

Access this article online

Quick Response Code:



Website:

www.jcritintensivecare.org

DOI:

10.14744/dcybd.2026.10864

# Cranial Nerve Involvement and Thalamic Lesions in West Nile Virus Encephalitis in a Kidney Transplant Recipient: Implications for Diagnosis and Prevention

Faruk Yildiz,<sup>1</sup> Volkan Inal<sup>2</sup>

To the Editor,

We read with great interest the case report by Gorgulu et al.,<sup>[1]</sup> titled “West Nile Virus Encephalitis in a Kidney Transplant Patient,” published in the *Journal of Critical and Intensive Care* 2025;16(3):138–141. The authors describe a rare case of West Nile virus (WNV) encephalitis in an immunosuppressed kidney transplant recipient who presented with fever, altered mental status, and progressive neurological deterioration. This report contributes significantly to the Turkish intensive care literature by illustrating the diagnostic challenges in immunosuppressed patients and the high mortality risk of neuroinvasive WNV disease.<sup>[1]</sup>

The most notable feature of this case is the simultaneous presence of bilateral thalamic lesions and contrast enhancement in the cisternal segments of the right trigeminal nerve and bilateral oculomotor nerves on MRI. While thalamic and basal ganglia hyperintensities are well recognized in WNV encephalitis, their combination with cranial nerve involvement—particularly of the trigeminal and oculomotor nerves—is rarely highlighted in the imaging litera-

ture. Although cranial nerve neuropathies (e.g., facial or abducens) have been reported in neuroinvasive WNV,<sup>[2]</sup> the specific pattern of cisternal enhancement involving the trigeminal and oculomotor nerves, alongside thalamic lesions, appears to be uncommon. In immunosuppressed hosts, this pattern may reflect meningoencephalitis with perineural spread and could serve as a highly suggestive radiological marker of WNV infection. This observation warrants validation in larger multicenter case series of immunocompromised patients.

Additionally, the occurrence of infection 36 months after transplantation further supports the possibility of donor-derived or transfusion-transmitted WNV.<sup>[3]</sup> Türkiye experienced its largest WNV outbreak in 2024, with 51 cases reported in a multicenter series and a case fatality rate of 17.6% (all fatal cases involved advanced age and comorbidities).<sup>[4]</sup> Given Türkiye’s endemic status and the increasing number of cases reported across Europe, routine WNV screening for organ donors and recipients remains nonstandard.<sup>[3,4]</sup> This case highlights the urgent need for a practical diagnostic algorithm: WNV PCR in cere-

<sup>1</sup>Department of Internal Medicine, Division of Intensive Care Medicine, Trakya University Faculty of Medicine, Edirne, Türkiye

<sup>2</sup>Department of Internal Medicine, Division of Intensive Care Medicine, Trakya University Faculty of Medicine, Edirne, Türkiye

**Address for correspondence:**

Faruk Yildiz, Department of Internal Medicine, Division of Intensive Care Medicine, Trakya University Faculty of Medicine, Edirne, Türkiye.

E-mail:

faruk\_yildiz@hotmail.com

**Received:** 03.04.2026

**Accepted:** 16.04.2026

**Published:** 20.04.2026

**How to cite this article:** Yildiz F, Inal V. Cranial Nerve Involvement and Thalamic Lesions in West Nile Virus Encephalitis in a Kidney Transplant Recipient: Implications for Diagnosis and Prevention. *J Crit Intensive Care* 2026;17(1):45–46.

This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License.

**For reprints contact:** kare@karepb.com

brospinal fluid should be routinely included in the meningoencephalitis panel for all immunosuppressed patients presenting with fever and altered consciousness, irrespective of season, and should be accompanied by prompt reduction of immunosuppression when indicated.<sup>[5]</sup>

We commend the authors for this insightful and unusual case report. Their findings emphasize the importance of reviewing WNV screening protocols in transplant centers and raising awareness of atypical radiological patterns in neuroinvasive viral infections among immunosuppressed patients. Multicenter prospective studies are necessary to enhance both diagnostic and preventive strategies.

**Conflict of Interest:** The authors have no conflicts of interest to declare.

**Funding:** The authors declared that this study received no financial support.

**Use of AI for Writing Assistance:** No use of AI-assisted technologies was declared by the authors.

**Author Contributions:** Concept – F.Y., V.I.; Design – F.Y., V.I.;

Supervision – V.I.; Data Collection and/or Processing - F.Y.; Analysis and/or Interpretation - V.I.; Literature Review – F.Y.; Writing – F.Y.; Critical Review – F.Y., V.

**Peer-review:** Internally peer-reviewed.

## References

1. Gorgulu RO, Usta O, Senturk YE, Ucem S, Ozserezli B, Sogut MS, et al. West Nile Virus Encephalitis in a Kidney Transplant Patient. *J Crit Intensive Care* 2025;16(3):138-141. [\[CrossRef\]](#)
2. Marshall EM, Bauer L, Nelemans T, Sooksawasdi Na Ayudhya S, Benavides F, et al. Differential susceptibility of human motor neurons to infection with Usutu and West Nile virus. *J Neuroinflammation* 2024;21(1):236. [\[CrossRef\]](#)
3. Abbas A, Qiu F, Sikyta A, Fey PD, Florescu DF. Neuroinvasive West Nile virus infections after solid organ transplantation: Single center experience and systematic review. *Transpl Infect Dis* 2022;24(6):e13929. [\[CrossRef\]](#)
4. Alhan Ö, Koç MM, Batrel A, Yılmaz E, Ersöz G, Aydın M, et al. Emerging West Nile virus infections in Türkiye. *Eur J Clin Microbiol Infect Dis* 2025;44(7):1659-66. [\[CrossRef\]](#)
5. Anesi JA, Silveira FP; AST Infectious Diseases Community of Practice. Arenaviruses and West Nile Virus in solid organ transplant recipients: Guidelines from the American Society of Transplantation Infectious Diseases Community of Practice. *Clin Transplant* 2019;33(9):e13576. [\[CrossRef\]](#)